

Alberta-Thy-22-97

hep-ph/9710275

August, 1997

Inelastic Final-State Interactions and Two-body Hadronic B decays into Single-Isospin channels

A.N.Kamal and C.W.Luo

*Theoretical Physics Institute and Department of Physics,
University of Alberta, Edmonton, Alberta T6G 2J1, Canada.*

Abstract

The role of inelastic final-state interactions in CP asymmetries and branching ratios is investigated in certain chosen single isospin two-body hadronic B decays. Treating final-state interactions through Pomeron and Regge exchanges, we demonstrate that inelastic final state interactions could lead to sizeable effects on the CP asymmetry.

PACS number(s): 11.30.Er, 13.25.Hw, 13.85.Fb

I. Introduction

It is well-known that CP asymmetries occur in two-body hadronic decays of B meson involving two distinct CKM angles and two differing strong phases[1]. The sources of the strong phases are several: perturbative penguin loops, final-state interaction (fsi) phases involving two different isospins, and inelastic final-state interactions involving a single isospin state.

In this paper we have studied the effect of interchannel mixing on B decays into two-body single-isospin channels. For reasons to follow, we have chosen to study the following decay modes: $B^- \rightarrow \eta_c \pi^-$, $B^- \rightarrow \eta_c K^-$ and $B^- \rightarrow \phi K^-$. The first of them, $B^- \rightarrow \eta_c \pi^-$, is a color-suppressed decay into a state with $I = 1$. At the tree-level, it proceeds through a CKM angle product $V_{cb}V_{cd}^*$. In absence of interchannel mixing, the CP asymmetry for this mode is known to vanish[2].(Though ref.[2] does not include electromagnetic penguins, their inclusion does not alter this fact.) The color-favored decay channel $B^- \rightarrow D^0 D^-$ with $I = 1$, also proceeds through a CKM product $V_{cb}V_{cd}^*$ at the tree-level, and has a nonvanishing CP asymmetry[2]. An inelastic coupling of $D^0 D^-$ channel to $\eta_c \pi^-$ allows a two-step decay $B^- \rightarrow D^0 D^- \rightarrow \eta_c \pi^-$ resulting in, as we show later, a significant CP asymmetry in $\eta_c \pi^-$ final state. The same can be argued for $B^- \rightarrow \eta_c K^-$, also a color-suppressed decay proceeding via $V_{cb}V_{cs}^*$ at the tree-level. It is also known to have a vanishing CP asymmetry[2] in absence of inelastic fsi. However, the interchannel mixing of $\eta_c K^-$ channel with $D^0 D_s^-$, a color-favored channel, results in a nonvanishing CP asymmetry in $\eta_c K^-$ mode. Lastly, the mode $B^- \rightarrow \phi K^-$ involves $b \rightarrow s \bar{s} s$ decay and proceeds only through a penguin amplitude involving CKM angle products $V_{ub}V_{us}^*$ and $V_{cb}V_{cs}^*$. In absence of inelastic fsi, it is known to have a nonzero CP asymmetry. The strong phases here arise from the light quark

penguin loops. However, ϕK^- channel can couple to $D^{*0}D_s^-$ and $D^0D_s^{*-}$ channels. The decays $B^- \rightarrow D^{*0}D_s^-$ and $B^- \rightarrow D^0D_s^{*-}$ being Cabibbo-favored, have branching ratios three orders of magnitude larger than that of the penguin process $B^- \rightarrow \phi K^-$ [2]. We have studied the effect of the inelastic coupling of the channels $B^- \rightarrow D^{*0}D_s^-$ and $D^0D_s^{*-}$ to $B^- \rightarrow \phi K^-$ channel on the branching ratio and CP asymmetry in the latter mode.

This paper is organized as follows: In section II, we describe the formalism and investigate the effect of inelastic fsi on the branching ratios and CP asymmetries in $B^- \rightarrow \eta_c \pi^-$, $B^- \rightarrow \eta_c K^-$ and $B^- \rightarrow \phi K^-$. The results are discussed in section III, .

II. CP Asymmetry and Final State Interaction in

$$B^- \rightarrow \eta_c \pi^-, B^- \rightarrow \eta_c K^- \text{ and } B^- \rightarrow \phi K^-$$

A. Definitions and Formalism

The effective Hamiltonian for $b \rightarrow s$ transition (for $b \rightarrow d$, replace s by d) is given by [3, 4, 5]

$$H_{eff} = \frac{G_F}{\sqrt{2}} \sum_{q=u,c} \left\{ V_{qb} V_{qs}^* [C_1 O_1^q + C_2 O_2^q + \sum_{i=3}^{10} C_i O_i] \right\}. \quad (1)$$

The operators in Eq.(1) are the following;

$$\begin{aligned} O_1^q &= (\bar{s}q)_{V-A}(\bar{q}b)_{V-A}, & O_2^q &= (\bar{s}_\alpha q_\beta)_{V-A}(\bar{q}_\beta b_\alpha)_{V-A}; \\ O_3 &= (\bar{s}b)_{V-A} \sum_{q'} (\bar{q}' q')_{V-A}, & O_4 &= (\bar{s}_\alpha b_\beta)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\alpha)_{V-A}, \\ O_5 &= (\bar{s}b)_{V-A} \sum_{q'} (\bar{q}' q')_{V+A}, & O_6 &= (\bar{s}_\alpha b_\beta)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\alpha)_{V+A}; \\ O_7 &= \frac{3}{2}(\bar{s}b)_{V-A} \sum_{q'} (e_{q'} \bar{q}' q')_{V+A}, & O_8 &= \frac{3}{2}(\bar{s}_\alpha b_\beta)_{V-A} \sum_{q'} (e_{q'} \bar{q}'_\beta q'_\alpha)_{V+A}, \\ O_9 &= \frac{3}{2}(\bar{s}b)_{V-A} \sum_{q'} (e_{q'} \bar{q}' q')_{V-A}, & O_{10} &= \frac{3}{2}(\bar{s}_\alpha b_\beta)_{V-A} \sum_{q'} (e_{q'} \bar{q}'_\beta q'_\alpha)_{V-A}. \end{aligned} \quad (2)$$

O_1 and O_2 are the Tree Operators, $O_3, \dots, O_6$ are generated by QCD Penguins and $O_7, \dots, O_{10}$ are generated by Electroweak Penguins. Here $V \pm A$ represent $\gamma_\mu(1 \pm \gamma_5)$, α and β are color indices. $\sum_{q'}$ is a sum over the active flavors u,d,s and c quarks.

In the next-to-leading-log calculation one works with effective Wilson coefficients C_i^{eff} , rather than the coefficients that appear in (1). The derivation of these effective coefficients is well known [3, 4, 5]. We simply quote their values

$$\begin{aligned} C_1^{eff} &= \bar{C}_1, & C_2^{eff} &= \bar{C}_2, & C_3^{eff} &= \bar{C}_3 - P_s/N_c, & C_4^{eff} &= \bar{C}_4 + P_s, \\ C_5^{eff} &= \bar{C}_5 - P_s/N_c, & C_6^{eff} &= \bar{C}_6 + P_s, & C_7^{eff} &= \bar{C}_7 + P_e, \\ C_8^{eff} &= \bar{C}_8, & C_9^{eff} &= \bar{C}_9 + P_e, & C_{10}^{eff} &= \bar{C}_{10}, \end{aligned} \quad (3)$$

with[3]

$$\begin{aligned} \bar{C}_1 &= 1.1502, & \bar{C}_2 &= -0.3125, & \bar{C}_3 &= 0.0174, & \bar{C}_4 &= -0.0373, & \bar{C}_5 &= 0.0104, & \bar{C}_6 &= -0.0459, \\ \bar{C}_7 &= -1.050 \times 10^{-5}, & \bar{C}_8 &= 3.839 \times 10^{-4}, & \bar{C}_9 &= -0.0101, & \bar{C}_{10} &= 1.959 \times 10^{-3}, \end{aligned} \quad (4)$$

and

$$P_s = \frac{\alpha_s(\mu)}{8\pi} C_1(\mu) \left[\frac{10}{9} + \frac{2}{3} \ln \frac{m_q^2}{\mu^2} - G(m_q, \mu, q^2) \right], \quad (5)$$

$$P_e = \frac{\alpha_{em}(\mu)}{3\pi} \left[C_2(\mu) + \frac{C_1(\mu)}{N_c} \right] \left[\frac{10}{9} + \frac{2}{3} \log \frac{m_q^2}{\mu^2} - G(m_q, \mu, q^2) \right], \quad (6)$$

where

$$G(m_q, \mu, q^2) = -4 \int_0^1 dx x(1-x) \ln \left[1 - x(1-x) \frac{q^2}{m_q^2} \right], \quad (7)$$

q^2 is the momentum carried by the gluon or the photon in the penguin diagram and m_q the mass of the quark q in the penguin loop. For $q^2 > 4m_q^2$, $G(m_q, \mu, q^2)$ becomes complex giving rise to strong perturbative phases through P_s and P_e . The parameters

we employ are:

$$m_u = 5MeV, m_s = 175MeV, m_c = 1.35GeV, m_b = 5.0GeV,$$

$$CKMangles : A = 0.81, \lambda = 0.22, (\rho, \eta) = (-0.20, 0.45) \text{ and } (0.30, 0.42),$$

$$f_D = 200MeV, f_{D_s} = f_{D_s^*} = f_{\eta_c} = 300MeV, f_\phi = 233MeV. \quad (8)$$

$$(9)$$

Consider now each one of the decays $B^- \rightarrow \eta_c \pi^-, \eta_c K^-$ and ϕK^- in absence of inelastic fsi. In the factorization approximation, which we adopt, the decay amplitudes are:

$$A(B^- \rightarrow \eta_c \pi^-) = \frac{G_F}{\sqrt{2}} \{V_{cb}V_{cd}^* a_2 - V_{tb}V_{td}^* (a_3 - a_5 - a_7 + a_9)\} \langle \eta_c | (\bar{c}c)_{V-A} | 0 \rangle \langle \pi^- | (\bar{d}b)_{V-A} | B^- \rangle \quad (10)$$

$$A(B^- \rightarrow \eta_c K^-) = \frac{G_F}{\sqrt{2}} \{V_{cb}V_{cs}^* a_2 - V_{tb}V_{ts}^* (a_3 - a_5 - a_7 + a_9)\} \langle \eta_c | (\bar{c}c)_{V-A} | 0 \rangle \langle K^- | (\bar{s}b)_{V-A} | B^- \rangle \quad (11)$$

$$A(B^- \rightarrow \phi K^-) = -\frac{G_F}{\sqrt{2}} V_{tb}V_{ts}^* \left\{ \left(1 + \frac{1}{N_c}\right) C_3^{eff} + \left(1 + \frac{1}{N_c}\right) C_4^{eff} + a_5 - a_7/2 - \frac{1}{2} \left(1 + \frac{1}{N_c}\right) C_9^{eff} \right. \\ \left. - \frac{1}{2} \left(1 + \frac{1}{N_c}\right) C_{10}^{eff} \right\} \langle \phi | (\bar{s}s)_{V-A} | 0 \rangle \langle K^- | (\bar{s}b)_{V-A} | B^- \rangle \quad (12)$$

where we have used the unitarity relation $\sum_{q=u,c} V_{qb}V_{qs(d)}^* = -V_{tb}V_{ts(d)}^*$ and defined

$$a_{2i} = C_{2i}^{eff} + \frac{1}{N_c} C_{2i-1}^{eff}, \\ a_{2i-1} = C_{2i-1}^{eff} + \frac{1}{N_c} C_{2i}^{eff}, \quad (13)$$

with i an integer. The strong phases appear through P_s and P_e defined in (5) and (6). However, in (10) and (11), odd coefficients a_3, a_5, a_7 and a_9 involve such combinations of C_i^{eff} as to cancel the effect of P_s and P_e . Thus, strong phases do not appear in (10) and (11) but they do in (12). Hence, CP asymmetry vanishes for $B^- \rightarrow \eta_c \pi^-$ and

$\eta_c K^-$ but is nonzero for $B^- \rightarrow \phi K^-$. This is true even if the electromagnetic penguins were ignored as in ref[2].

Let us now consider the decay channel $B^- \rightarrow D^0 D^-$. The decay amplitude is given by,

$$A(B^- \rightarrow D^0 D^-) = \frac{G_F}{\sqrt{2}} \{V_{cb}V_{cd}^* a_1 - V_{tb}V_{td}^* (a_4 + (a_6 + a_8)R_1 + a_{10})\} \langle D^- | (\bar{d}c)_{V-A} | 0 \rangle \langle D^0 | (\bar{c}b)_{V-A} | B^- \rangle \quad (14)$$

where

$$R_1 = \frac{2m_D^2}{(m_b - m_c)(m_c + m_d)}. \quad (15)$$

Note that the above decay is color-favored (tree diagram being proportional to a_1) and that strong phases do not cancel in the even coefficients a_4, a_6, a_8 and a_{10} . Hence CP asymmetry in $B^- \rightarrow D^0 D^-$ is nonvanishing[2]. Throughout our calculations, we have used the formfactors from Bauer, Stech and Wirbel[6].

In the following section we discuss in detail the mixing of $D^0 D^-$ and $\eta_c \pi^-$ channels through inelastic fsi.

B. Inelastic mixing of $D^0 D^-$ and $\eta_c \pi^-$ channels.

Inelastic fsi have been discussed in the past[7, 8, 9, 10, 11] in the context of the K-matrix formalism. The desirable feature of this method is that unitarity of the S-matrix is ensured. The difficulty lies in the proliferation of K-matrix parameters, mostly unknown, with the number of channels. Moreover, in two-channel problems, the second channel (the inelastic channel) is assumed to reflect (through unitarity) all the inelastic channels. This is an oversimplification of reality. In ref.[8], the coupling of $\eta_c \pi^-$ channel to $\eta \pi^-$ and $\eta' \pi^-$ is discussed in K-matrix formalism. We comment on this work in Section III.

In the calculations we present below, we make no effort to enforce two-channel unitarity. Rather, we couple the decay channel $\eta_c\pi^-$ to D^0D^- using a Regge-exchange model. The model coupling constants are related to known coupling constants by approximations we explain in the text. The advantage of this procedure, in contrast to the K-matrix approach[7, 8, 9, 10, 11], is that the scattering parameters are determined more realistically. The shortcoming is that the relevant elements of the S-matrix being completely determined, the S-matrix itself does not satisfy two-channel unitarity. Yet, we think it is more realistic to treat the effect of inelastic channels one channel at a time rather than enforce two-channel unitarity on what is in fact a multi-channel problem.

We begin by establishing certain key equations for an arbitrary number of two-body channels. An $n \times n$ S-matrix for s-wave scattering satisfying unitarity can be written as [12]

$$\mathbf{S} = (\mathbf{1} + i\mathbf{k}^{\frac{1}{2}}\mathbf{K}\mathbf{k}^{\frac{1}{2}})(\mathbf{1} - i\mathbf{k}^{\frac{1}{2}}\mathbf{K}\mathbf{k}^{\frac{1}{2}})^{-1}, \quad (16)$$

where $\mathbf{k}$ is a diagonal momentum matrix and $\mathbf{K}$ a real-symmetric matrix with $n(n+1)/2$ real parameters. The decay amplitudes for the B meson into n two-body channels are inelastically coupled through

$$\mathbf{A} = (\mathbf{1} - i\mathbf{k}^{\frac{1}{2}}\mathbf{K}\mathbf{k}^{\frac{1}{2}})^{-1}\mathbf{A}^{(0)}, \quad (17)$$

where $\mathbf{A}^{(0)}$ is a column of uncoupled amplitudes. The coupled (unitarized) amplitudes are assembled in the column $\mathbf{A}$. From (16) and (17) it is easily shown that [11]

$$\mathbf{A} = \frac{\mathbf{1} + \mathbf{S}}{2}\mathbf{A}^{(0)}. \quad (18)$$

Let us label channels $\eta_c\pi^-$ and D^0D^- as channels 1 and 2 respectively. $A_1^{(0)}$ and $A_2^{(0)}$ are given by (10) and (14) respectively. In order to calculate the effect of channel 2 on channel 1 and vice-versa, we need to calculate the elements S_{11} , S_{12} and S_{22} of the S-matrix . We describe their evaluation in the following.

We assume that Pomeron exchange dominates elastic scattering. The scatterings $\eta_c\pi^- \rightarrow \eta_c\pi^-$ and $D^0D^- \rightarrow D^0D^-$ are then represented by amplitudes of the form[13]

$$P(s, t) = \beta(t) \left(\frac{s}{s_0}\right)^{\alpha_P(t)} e^{i\pi\alpha_P(t)/2} \quad (19)$$

where $\sqrt{s_0}$ is an energy scale and the Pomeron trajectory is parameterized by

$$\alpha_P(t) = 1.08 + 0.25t. \quad (20)$$

The momentum transfer t is expressed in GeV^2 in the above. The Pomeron coupling strength, $\beta(t)$, is assumed[14, 15] to have a t -dependence of the form $\beta(t) = \beta(0)e^{2.8t}$. In the additive quark model, $\beta(0) = 4\beta(cu)$ for $\eta_c\pi^- \rightarrow \eta_c\pi^-$, and $\beta(0) = 2\beta(cu) + \beta(uu) + \beta(cc)$ for $D^0D^- \rightarrow D^0D^-$ scattering. The residue $\beta(uu)$ can be extracted from high energy pp and πp scattering data yielding [14, 15] $\beta(uu) \approx 6.5$. No experimental information exists for the determination of $\beta(cu)$. We make the theoretical ansatz[15]: $\beta(cu) \approx \frac{1}{10}\beta(uu)$, and assume $\beta(cc)$ to be negligibly small[15]

The inelastic scattering $\eta_c\pi^- \rightarrow D^0D^-$ is mediated by D^{*0} Regge-exchange in the t channel (defined by $t = (P_{D^0} - P_{\eta_c})^2$). The amplitude is of the form [16]

$$R(s, t) = \beta_R(t) \frac{1 - \exp[-i\pi\alpha(t)]}{\sin[\pi\alpha(t)]} \left(\frac{s}{s_0}\right)^{\alpha(t)}. \quad (21)$$

For $\beta_R(t)$ we adopt a t dependence[13, 16, 17]

$$\beta_R(t) = \frac{\beta_R(0)}{\Gamma[\alpha(t)]}. \quad (22)$$

The fact that $\Gamma(z)$ has simple poles at $z = 0, -1, -2, \dots$, ensures that the Regge amplitude (21) does not develop nonsense poles at $\alpha(t) = 0, -1, -2, \dots$. We also note that in addition to $R(s, t)$ of (21), there is a u -channel exchange amplitude $R(s, u)$ generated by a charged D^* exchange.

Generally, s_0 is expected to be process-dependent. For light mesons and baryons it has been taken[13] as $s_0 = \frac{1}{\alpha'} \approx 1GeV^2$. However, for heavy mesons and baryons, the scale s_0 must reflect somehow a higher threshold for the scattering processes. Based on the work of [17], it is argued in [16] that for πD scattering mediated by ρ -trajectory, $s_0^{\pi D} \approx \frac{2}{\alpha'_R}$. We assume this value for $\eta_c \pi^- \rightarrow D^0 D^-$ scattering amplitude.

We determine $\beta_R(0)$ by taking the limit $\alpha(t) \rightarrow 1$ (D^* pole) in (21) and comparing it with the perturbative t-channel D^* -pole diagram. For the latter we assume a VPP vertex of form:

$$L_{int}^{VPP} = f_{ijk} g_{VPP} V_\mu^i P^j \partial^\mu P^k, \quad (23)$$

where i, j and k are SU(4) labels and f_{ijk} the antisymmetric symbol.

The perturbative t-channel D^* -exchange graph yields,

$$R(s \rightarrow \infty, t \sim m_{D^*}^2) = g_{VPP}^2 \frac{s}{t - m_{D^*}^2}. \quad (24)$$

Comparing the limiting case ($t \rightarrow m_{D^*}^2, \alpha(m_{D^*}^2) = 1$) of (21) with (24) results in

$$\beta_R(0) = \pi g_{VPP}^2. \quad (25)$$

SU(4) symmetry allows us to determine g_{VPP} from $\rho \rightarrow \pi\pi$ and $K^* \rightarrow K\pi$ decays[18],

$$\frac{g_{VPP}^2}{4\pi} \approx 3.0. \quad (26)$$

Heavy Quark effective Theory(HQET)[19, 20, 21, 22] could also have been used to determine $g_{D^* D \pi}$ if the rate $\Gamma(D^* \rightarrow D\pi)$ were known. In absence of this information, authors of ref.[19] fix this coupling by constraining it to yield the axial coupling of the nucleon, $g_A \approx 1.25$. This results in

$$g_{D^*-D^0\pi^-} = g_{D^*0D^+\pi^-} = g_{D_s^{*0}D^+K^-} \approx \frac{3\sqrt{m_D m_{D^*}}}{4f_\pi} \quad to \quad \frac{\sqrt{m_D m_{D^*}}}{f_\pi}, \quad (27)$$

where $f_\pi = 131 MeV$. This is a much larger coupling constant than that implied in [18], resulting in $\Gamma(D^{*-} \rightarrow D^0 \pi^-) = (100 - 180) KeV$ [19]. In contrast, the SU(4) symmetry scheme of ref[18] obtains $\Gamma(D^{*-} \rightarrow D^0 \pi^-) = 16 KeV$ using g_{VPP} given in (26). In our calculations we use the SU(4) symmetry coupling given by (23) and (26) only.

From the scattering amplitudes we project out the elements of the S-wave scattering matrix

$$S_{ij} = \delta_{ij} + \frac{i}{8\pi\sqrt{\lambda_i\lambda_j}} \int_{t_{min}}^{t_{max}} dt T(s, t), \quad (28)$$

where $T(s, t)$ is the total amplitude, λ_i and λ_j are the usual triangle functions $\lambda(x, y, z) = (x^2 + y^2 + z^2 - 2xy - 2xz - 2yz)^{1/2}$ for channels i and j respectively and t_{max}, t_{min} are the limits of the momentum transfer. We also took into account the u-channel charged D^* -exchange in calculating the S-matrix elements. The resulting S-matrix elements (channel 1 = $\eta_c \pi^-$, channel 2 = $D^0 D^-$) are,

$$S = \begin{pmatrix} 0.946 - 0.93 \times 10^{-3}i & 0.19 \times 10^{-2} + 0.068i & \\ 0.19 \times 10^{-2} + 0.068i & 0.843 - 0.27 \times 10^{-2}i & \\ . & . & \end{pmatrix}. \quad (29)$$

Clearly, two-channel unitarity is not satisfied but the S-matrix elements $S_{ij}(i, j = 1, 2)$ are completely determined. Calculation of the unitarized decay amplitudes proceeds by using $A_i^{(0)}$ from (10) and (14), and $\mathbf{S}$ from (28) in (18). The calculation of the branching ratios and CP asymmetry is then straight forward. We have chosen to perform the calculation for $N_c = 3$ and $N_c = 2.4$. The latter choice, suggested in [23], could be interpreted to reflect nonfactorization effects. The results are shown in Tables 1 and 2.

We note from these Tables that the induced CP asymmetry in $\eta_c \pi$ channel is large; in fact, as large as in channel $D^0 D^-$ to which it is coupled. The CP asymmetry in $\eta_c \pi^-$ channel, however, depends almost linearly on g_{VPP}^2 . Thus, increasing (decreasing) g_{VPP}

by a factor of 2 results in an increase (decrease) of CP asymmetry by approximately a factor of four. We defer the discussion of the results to Section III.

C. Inelastic Mixing of $D^0 D_s^-$ and $\eta_c K^-$ channels

In absence of interchannel coupling, the decay amplitude for $B^- \rightarrow D^0 D_s^-$ is given by

$$A(B^- \rightarrow D^0 D^-) = \frac{G_F}{\sqrt{2}} \{V_{cb}V_{cs}^* a_1 - V_{tb}V_{ts}^* (a_4 + (a_6 + a_8)R_2 + a_{10})\} \langle D_s^- | (\bar{s}c)_{V-A} | 0 \rangle \langle D^0 | (\bar{c}b)_{V-A} | B^- \rangle, \quad (30)$$

where

$$R_2 = \frac{2m_{D_s}^2}{(m_b - m_c)(m_c + m_s)}. \quad (31)$$

Let us label $\eta_c K^-$ and $D^0 D_s^-$ as channels 1 and 2 respectively. The pomeron-mediated elastic scattering now involves coupling constants $2\beta(cu) + 2\beta(cs)$ for $\eta_c K^-$ channel and $\beta(cc) + \beta(cu) + \beta(cs) + \beta(us)$ for $D^0 D_s^-$ channel. For $\beta(cs)$ we use the ansatz: $\beta(cs) \approx \frac{1}{10}\beta(us) \approx \frac{1}{15}\beta(uu)$. The Pomeron amplitude is then given as in (19).

The inelastic scattering $\eta_c K^- \rightarrow D^0 D_s^-$ is mediated by D^{*0} -exchange in the t channel ($t = (P_{D^0} - P_{\eta_c})^2$), and by D_s^* -exchange in the u channel. The calculation of the effect of inelastic coupling of $\eta_c K^-$ and $D^0 D_s^-$ channels parallels that of $\eta_c \pi^-$ and $D^0 D^-$ channels described in the previous section. The resulting S-matrix ($\eta_c K^- =$ channel 1, $D^0 D_s^- =$ channel 2) is

$$S = \begin{pmatrix} 0.954 - 0.8 \times 10^{-3}i & 0.85 \times 10^{-3} + 0.068i & \dots \\ 0.85 \times 10^{-3} + 0.068i & 0.888 - 0.19 \times 10^{-2}i & \dots \\ . & . & \dots \end{pmatrix}. \quad (32)$$

The resulting branching ratios and CP asymmetries are shown in Tables 1 and 2. Again, we notice that CP asymmetry induced in channel $\eta_c K^-$ is comparable to that in channel $D^0 D_s^-$. Further discussion of the results is deferred to section III.

D. Inelastic coupling of ϕK^- to $D^{*0} D_s^-$ and $D_s^{*-} D^0$ channels.

In absence of inelastic fsi, the decay amplitudes for $B^- \rightarrow D^{*0} D_s^-$ and $D_s^{*-} D^0$ are:

$$\begin{aligned} A(B^- \rightarrow D^{*0} D_s^-) &= \frac{G_F}{\sqrt{2}} \{V_{cb} V_{cs}^* a_1 - V_{tb} V_{ts}^* (a_4 - (a_6 + a_8) R_3 + a_{10})\} \langle D_s^- | (\bar{s}c)_{V-A} | 0 \rangle \langle D^{*0} | (\bar{c}b)_{V-A} | B^- \rangle \\ A(B^- \rightarrow D^0 D_s^{*-}) &= \frac{G_F}{\sqrt{2}} \{V_{cb} V_{cs}^* a_1 - V_{tb} V_{ts}^* (a_4 + a_{10})\} \langle D_s^{*-} | (\bar{s}c)_{V-A} | 0 \rangle \langle D^0 | (\bar{c}b)_{V-A} | B^- \rangle, \end{aligned}$$

where

$$R_3 = \frac{2m_{D_s}^2}{(m_s + m_c)(m_c + m_b)}. \quad (35)$$

Inelastic fsi couple the amplitude for $B^- \rightarrow \phi K^-$, eq.(12), to the amplitudes in (33) and (34). The calculation of the S-matrix elements (ϕK^- = channel 1, $D_s^- D^{*0}$ = channel 2, $D_s^{*-} D^0$ = channel 3) is considerably simplified in the B rest-frame. This is because in this frame the vector meson can only have longitudinal helicity in a $B \rightarrow VP$ decay. Because of this fact, the Pomeron and Regge amplitudes involving only helicity 0 $\rightarrow$ helicity 0 transition are the same as in spin-less scattering. The Pomeron amplitude for elastic scattering is given by (19) with $\beta(0) = 2\beta(su) + 2\beta(ss)$ for ϕK^- elastic scattering, and $\beta(0) = \beta(us) + \beta(cu) + \beta(cs) + \beta(cc)$ for $D_s^- D^{*0}$ and $D_s^{*-} D^0$ elastic scatterings. We assume $\beta(ss) \approx \frac{2}{3}\beta(su)$.

Inelastic scatterings, $\phi K^- \rightarrow D_s^- D^{*0}$ and $D_s^{*-} D^0$, are mediated by D_s^* exchange in the t channel ($t = (P_{D_s} - P_\phi)^2$ and $(P_{D_s^*} - P_\phi)^2$ respectively). There are no u channel exchanges. D_s -trajectory, being a lower-lying trajectory, makes a smaller contribution and we neglect it. The Regge-amplitude is assumed to be of the form given in (21). In order to determine the coupling $\beta(0)$, we equate the limiting form of (21) for $t \rightarrow m_{D_s^*}^2$ with the perturbative expressions for $\phi K^- \rightarrow D_s^- D^{*0}$ and $D_s^{*-} D^0$ with D_s^* exchange. We adopt the following definitions,

$$D_s^* - trajectory : \quad \alpha_{D_s^*} \approx -1.23 + 0.5t, \quad (36)$$

$$VVP - vertex : \quad L_{int}(VVP) = g_{VVP} d_{ijk} \epsilon_{\mu\nu\rho\sigma} \partial_\mu V_\nu^i \partial_\rho V_\sigma^j P^k, \quad (37)$$

$$VVV - vertex : \quad L_{int}(VVV) = g_{VVV} f_{ijk} (\partial_\mu V_\nu^i - \partial_\nu V_\mu^i) V_\mu^j V_\nu^k, \quad (38)$$

where d_{ijk} and f_{ijk} are SU(4) indices. The coupling g_{VVP} has dimension $(mass)^{-1}$ while g_{VVV} is dimensionless. D_s^* trajectory is assumed to be parallel to D^* trajectory with a slope as in[14].

The evaluation of perturbative D_s^* -exchange diagram was done numerically. The calculation was made simpler by the fact that the vector particles could only be longitudinally polarized in B rest-frame. For large s we obtain for the digrams shown in Figs.1 and 2,

$$\begin{aligned} T^{Fig.1}(s \rightarrow \infty, t \sim m_{D_s^*}^2) &= 0.7 GeV^2 g_{D^* D_s^* K} g_{D_s D_s^* \phi} \frac{s}{t - m_{D_s^*}^2}, \\ T^{Fig.2}(s \rightarrow \infty, t \sim m_{D_s^*}^2) &= 0.5 g_{D D_s^* K} g_{D_s^* D_s^* \phi} \frac{s}{t - m_{D_s^*}^2}. \end{aligned} \quad (39)$$

The corresponding Regge-exchange amplitude yields

$$T(s \rightarrow \infty, t \sim m_{D_s^*}^2) = \frac{\beta(0)}{\pi} \frac{s}{t - m_{D_s^*}^2}, \quad (40)$$

where we have used $s_0 = 2/\alpha'$.

A comparison of (40) with (39) yields $\beta(0)$. The coupling constants in (39) are determined as follows: In HQET[19], where light pseudoscalar mesons are introduced as nonlinear realization of $SU(3)_L \times SU(3)_R$, one obtains ($f_\pi = 131 MeV$, and we are using the parameter[19] $f=-1.5$),

$$g_{D^{*0} D_s^{*-} K^-} \approx \frac{3}{4f_\pi}. \quad (41)$$

Light vector and axial-vector mesons can also be introduced in HQET[20, 21, 22] allowing a $D_s^* D_s \phi$ coupling. However, HQET does not by itself permit an evaluation of this coupling constant. Use of vector dominance in the radiative decays of light mesons determines [24] $g_{VVP} \approx 6\text{GeV}^{-1}$. Using SU(4) symmetry, one obtains $g_{D_s^{*-} D_s^- \phi} \approx 6/\sqrt{2}\text{GeV}^{-1}$ which is a little lower than the value given in (41). For want of a better choice we assume $g_{D_s^{*-} D^- \phi} = g_{D^{*0} D_s^{*-} K^-}$.

HQET[20, 21, 22] allows us to calculate the VVV coupling $g_{D_s^* D_s^* \phi}$ provided an assumption is made as to how the flavor singlet and the flavor octet of the light vector meson couple to $D_s^* D_s^*$. In nonet symmetry, we obtain

$$g_{D_s^* D_s^* \omega} = 0 \quad (42)$$

$$g_{D_s^* D_s^* \phi} = g_{VVV} = g \text{ (of [22]) } \approx 4.3 \quad (43)$$

As for the VPP coupling, we adopt the value in (26).

To calculate the effect of channels $D^{*0} D_s^-$ (channel 2) and $D_s^{*-} D^0$ (channel 3) on ϕK^- (channel 1), we need the elements S_{11}, S_{12}, S_{13} of the S-matrix. The decay amplitude for $B^- \rightarrow \phi K^-$ is then given by,

$$A(B^- \rightarrow \phi K^-) = \frac{1 + S_{11}}{2} A^{(0)}(B^- \rightarrow \phi K^-) + \frac{S_{12}}{2} A^{(0)}(B^- \rightarrow D^{*0} D_s^-) + \frac{S_{13}}{2} A^{(0)}(B^- \rightarrow D^0 D_s^{*-}), \quad (44)$$

The relevant elements of the S-matrix are calculated to be,

$$\begin{aligned} S_{11} &= 0.784 - 0.37 \times 10^{-2}i \\ S_{12} &= -0.53 \times 10^{-3} + 0.12 \times 10^{-2}i \\ S_{13} &= -0.33 \times 10^{-3} + 0.70 \times 10^{-3}i \end{aligned} \quad (45)$$

Since $B^- \rightarrow \phi K^-$ is a penguin mediated process, its branching ratio in absence of inelastic coupling is small ($\sim 10^{-6}$ to 10^{-5})[2]. In contrast, the inelastic channels

$D^{*0}D_s^-$ and $D^0D_s^{*-}$ to which it couples are Cabibbo-favored and have branching ratios of the order of 10^{-3} to 10^{-2} [2]. Thus, whereas the two channels $D^{*0}D_s^-$ and $D^0D_s^{*-}$ can influence the branching ratios and CP asymmetries in ϕK^- , we do not expect the ϕK^- channel to significantly effect the channels $D^{*0}D_s^-$ and $D^0D_s^{*-}$. Further, the inelastic coupling of $D^{*0}D_s^-$ channel to $D^0D_s^{*-}$ occurs via the exchange of $(c\bar{c})$ mesonic trajectories, η_c and ψ . As both of these trajectories are low-lying with large and negative intercepts $\alpha(0)$, their contribution to the inelastic scattering $D^{*0}D_s^- \rightarrow D^0D_s^{*-}$ is expected to be highly suppressed. We, therefore, do not expect one of these channels to effect the other significantly either. For these reasons we have displayed in Tables 1 and 2 only the effect on the channel ϕK^- . The other two channels, $D^{*0}D_s^-$ and $D^0D_s^{*-}$, are left largely unaffected by fsi.

Tables 1 and 2 show that though the effect of inelastic fsi on the branching ratio for ϕK^- is small (due to the small size of S_{12} and S_{13} in (45)), the effect on the CP asymmetry is significant. The results are discussed in the following section.

III. Results and Discussion

In absence of inelastic fsi, CP asymmetries in $B^- \rightarrow \eta_c \pi^-$ and $\eta_c K^-$ channels vanish[2]. We have shown that an inelastic coupling of $\eta_c \pi^-$ channel to $D^0 D^-$ and that of $\eta_c K^-$ to $D^0 D_s^-$, leads to substantial CP asymmetries in $B^- \rightarrow \eta_c \pi^-$ and $\eta_c K^-$ decays. We have used $N_c = 3$ and 2.4 and two sets of values for (ρ, η) . The CP asymmetries depend sensitively on the coupling constant g_{VPP} and the energy-scale parameter s_0 . An increase (decrease) of g_{VPP} by a factor of two increases (decreases) the CP asymmetry by roughly a factor of 4. Similarly, increasing s_0 from $1 GeV^2$ to $2/\alpha'_R$ enhances the CP asymmetry by raising the values of the off-diagonal elements of the S-matrix. In addition, the calculated CP asymmetry will also depend on the

effective q^2 employed. We have used $q^2 = m_b^2/2$.

Ref.[8] discusses a three-channel problem, $B^- \rightarrow \eta\pi^-, \eta'\pi^-$ and $\eta_c\pi^-$, in the K-matrix formalism and demonstrates that a CP asymmetry of the order of $\sim 1\%$ can be generated in $\eta_c\pi^-$ channel through inelastic fsi. Our work differs from [8] in several respects. Most importantly, we couple $\eta_c\pi^-$, a color-suppressed channel, to D^0D^- , a color-favored channel. The channels, $\eta\pi^-$ and $\eta'\pi^-$, invoked in [8] are both color-suppressed, and thus, not expected to be as important as D^0D^- . In addition, we do not require the S-matrix to be unitary at two or three channel level. Another important difference between our work and that of [8] lies in how the strong interaction fsi parameters are determined. We use Pomeron and Regge phenomenology, presumably applicable at $\sqrt{s} \sim m_B$, while [8] determines the K-matrix elements through low-energy phenomenology. For example, the diagonal and off-diagonal elements of the K-matrix are evaluated using a contact ϕ^4 -interaction which allows for only S-wave scattering. This is expected to be a reasonable approximation at threshold but hardly likely to hold at $\sqrt{s} \sim m_B$. Despite these differences, we emphasize that the important conclusion of [8] was that a significant CP asymmetry in $\eta_c\pi^-$ could be generated by coupling it to $\eta\pi^-$ and $\eta'\pi^-$ channels. However, the fact that they also found asymmetries of the order of 10% and 20% in $\eta\pi^-$ and $\eta'\pi^-$ channels has little to do with inelastic fsi; asymmetries of this magnitude are generated in these channels in absence of inelastic fsi[2].

We have found that the CP asymmetry in $B^- \rightarrow \phi K^-$ (a penguin driven process) is significantly effected by a coupling to Cabibbo-favored channels $D^{*0}D_s^-$ and $D^0D_s^{*-}$. Due to the smallness of the off-diagonal elements of the S-matrix, the effect on the branching ratio is not as large as on CP asymmetry. Again, the CP asymmetry depends sensitively on the coupling constants and the value of s_0 .

This work was partly supported by a research grant from the Natural Sciences and Engineering Research Council of Canada to A.N.Kamal.

Table 1: Branching ratios and CP asymmetries with $q^2 = m_b^2/2$ and $N_c = 3$ (entries in bracket correspond to the uncoupled case)

CKM Matrix	$\rho = 0.30, \quad \eta = 0.42$		$\rho = -0.20, \quad \eta = 0.45$	
Decay channel	BR	$a_{CP}(\%)$	BR	$a_{CP}(\%)$
$B^- \rightarrow \eta_c \pi^-$	2.85×10^{-6}	3.97	2.87×10^{-6}	4.24
	(2.32×10^{-6})	(0.00)	(2.33×10^{-6})	(0.00)
$B^- \rightarrow D^0 D^-$	3.55×10^{-4}	3.94	3.22×10^{-4}	4.69
	(4.18×10^{-4})	(3.97)	(3.80×10^{-4})	(4.72)
$B^- \rightarrow \eta_c K^-$	7.42×10^{-5}	-0.25	7.42×10^{-5}	-0.27
	(5.54×10^{-5})	(0.00)	(5.54×10^{-5})	(0.00)
$B^- \rightarrow D^0 D_s^-$	1.47×10^{-2}	-0.21	1.48×10^{-2}	-0.23
	(1.65×10^{-2})	(-0.21)	(1.66×10^{-2})	(-0.23)
$B^- \rightarrow \phi K^-$	6.21×10^{-6}	0.74	5.97×10^{-6}	0.83
	(7.75×10^{-6})	(0.58)	(7.44×10^{-6})	(0.65)

Table 2: Branching ratios and CP asymmetries with $q^2 = m_b^2/2$ and $N_c = 2.4$ (entries in bracket correspond to the uncoupled case)

CKM Matrix	$\rho = 0.30,$	$\eta = 0.42$	$\rho = -0.20,$	$\eta = 0.45$
Decay channel	BR	$a_{CP}(\%)$	BR	$a_{CP}(\%)$
$B^- \rightarrow \eta_c \pi^-$	1.31×10^{-5}	1.80	1.32×10^{-5}	1.90
	(1.29×10^{-5})	(0.00)	(1.29×10^{-5})	(0.00)
$B^- \rightarrow D^0 D^-$	3.38×10^{-4}	3.74	3.06×10^{-4}	4.45
	(3.98×10^{-4})	(3.80)	(3.61×10^{-4})	(4.52)
$B^- \rightarrow \eta_c K^-$	3.12×10^{-4}	-0.12	3.11×10^{-4}	-0.13
	(3.08×10^{-4})	(0.00)	(3.08×10^{-4})	(0.00)
$B^- \rightarrow D^0 D_s^-$	1.29×10^{-2}	-0.19	1.30×10^{-2}	-0.20
	(1.45×10^{-2})	(-0.19)	(1.46×10^{-2})	(-0.21)
$B^- \rightarrow \phi K^-$	7.36×10^{-6}	0.58	7.06×10^{-6}	0.65
	(9.14×10^{-6})	(0.43)	(8.75×10^{-6})	(0.48)

References

- [1] See, for example, G.Kramer, W.F.Palmer and H.Simma, Nucl.Phys.B428, 77(1994).
- [2] G.Kramer, W.F.Palmer and H.Simma, Z.Phys.C66, 429(1995).
- [3] N.G.Deshpande and X.G.He, Phys.Rev.Lett. 74, 26(1995); Phys.Lett.B336, 471(1994).
- [4] R.Fleischer, Z.Phys.C62, 81(1994).
- [5] A.J.Buras, M.Jamin, M.Lautenbacher and P.Weisz, Nucl.Phys.B400, 37 (1993); A.J.Buras, M.Jamin and M.Lautenbacher, ibid. 75 (1993); M.Ciuchini, E.Franco, G.Martinelli and L.Reina, Nucl.Phys.B415, 403(1994).
- [6] M.Bauer, B.Stech and M.Wirbel, Z.Phys.C34, 103(1987).
- [7] A.N.Kamal, N.Sinha and R.Sinha, Z.Phys.C41, 207(1988)
- [8] S.Barshay, D.Rein and L.M.Sehgal, Phys.Lett.B259,475(1991).
- [9] M.Wanninger and L.M.Sehgal, Z.Phys.C50, 47(1994).
- [10] A.N.Kamal, Int.J.Mod.Phys.A7, 3515(1992).
- [11] A.N.Kamal and C.W.Luo, Alberta-Thy-02-97, hep-ph/9702289 (unpublished).
- [12] R.G.Newton, *Scattering Theory of Waves and Particles*, Mc-Graw-Hill, N.Y.(1996).
- [13] P.D.B.Collins, *Introduction to Regge Theory and High Energy Physics* (Cambridge Univ.Press, 1977).

- [14] H.Q.Zheng, Phys.Lett.B356,107(1995).
- [15] G.Nardulli and T.N.Pham, Phys.Lett.B391, 165 (1997).
- [16] B.Blok and I.Halperin, Phys.Lett.B385, 324 (1996).
- [17] P.E.Volkovitsky and A.B.Kaidalov, Yad.Fiz.35,1231, 1556(1982).
- [18] R.L.Thews and A.N.Kamal, Phys.Rev.D32, 810(1985).
- [19] T.M.Yan, H.Y.Cheng, C.Y.Cheung, G.L.Lin, Y.C.Lin and H.L.Yu, Phys.Rev.D46, 1148(1992).
- [20] J.Schechter and A.Subbaraman, Phys.Rev.D48, 332(1993).
- [21] R.Casalbuoni, A.Deandrea, N.D.Bartolomeo, R.Gatto, F.Feruglio and G.Nardulli, Phys.Lett.B299, 139(1993).
- [22] A.N.Kamal and Q.P.Xu, Phys.Rev.D49, 1526(1994).
- [23] A.N.Kamal and T.N.Pham, Phys.Rev.D50, 395(1994).
- [24] B.J.Edwards, *Radiative Decay of Mesons*, Ph.D thesis(1978), University of Alberta

Figure Captions

Fig.1: Inelastic scattering $B^- \rightarrow D^{*0} D_s^- \rightarrow \phi K^-$ through D_s^* exchange

Fig.2: Inelastic scattering $B^- \rightarrow D^0 D_s^{*-} \rightarrow \phi K^-$ through D_s^* exchange




Fig.1




Fig.2